

Description	Precipitating polyclonal donkey antiserum to goat IgG, heavy and light chains																
Product code	DoAG/IgG(H+L)																
Biological origin	Donkey																
Physical form	Delipidated, heat inactivated lyophilized stable whole serum.																
Preservative	No preservative added, as it may interfere with the antibody activity. No foreign protein added.																
Immunogen	Purified normal IgG isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.																
Adsorption	No adsorption required.																
Identity & Specificity	The reactivity of the antiserum is directed to the Fc and Fab subunits of both subclasses, IgG1 and IgG2. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. In immunoelectrophoresis against goat serum a single characteristic precipitin line with a spur is usually obtained, representing primarily IgG2. If the IgG1 level in the sample approaches the concentration of IgG2, it may become visible as a second precipitin line. It does not react with any non-Ig protein in goat serum, as tested by immunoelectrophoresis and double radial immunodiffusion.																
Cross-reactivity	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has been tested in double radial immunodiffusion with the following results: <table><tr><td>bovine ++</td><td>guinea pig ±</td><td>mouse ±</td><td>rat ±</td></tr><tr><td>cat ±</td><td>hamster -</td><td>monkey +</td><td>sheep ++</td></tr><tr><td>chicken -</td><td>horse -</td><td>pigeon -</td><td>swine -</td></tr><tr><td>dog ±</td><td>human +</td><td>rabbit ±</td><td>turkey -</td></tr></table>	bovine ++	guinea pig ±	mouse ±	rat ±	cat ±	hamster -	monkey +	sheep ++	chicken -	horse -	pigeon -	swine -	dog ±	human +	rabbit ±	turkey -
bovine ++	guinea pig ±	mouse ±	rat ±														
cat ±	hamster -	monkey +	sheep ++														
chicken -	horse -	pigeon -	swine -														
dog ±	human +	rabbit ±	turkey -														
Protein concentration	Total protein and IgG concentration in the antiserum are comparable to those of pooled donkey serum. No foreign proteins added.																
Antibody titre	Precipitin titre not less than 1:128 when tested against normal goat serum in agar block titration.																
Intended use	In immunoelectrophoresis and double radial immunodiffusion to identify the presence of IgG; as secondary antibody to precipitate the immunoglobulin in normal goat serum; to prepare an immunoabsorbent for the production of Ig-free goat serum, plasma or non-Ig preparations. This antiserum is not intended for use in non-precipitating antibody-binding or other highly sensitive assays. For such uses labelled and unlabelled cytochemical and immunoassay reagents meeting the necessary additional specificity and performance requirements are available.																
Directions for use	In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.																
Handling	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.																
Packing	Vial with 1 ml lyophilized antiserum.																
Storage / shelf life	Lyophilized at +4°C at least 10 years reconstituted at or below -20°C 3-5 years reconstituted at +4°C 7 days																
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.																

NORDIC IMMUNOLOGICAL LABORATORIES

Langendijk 25, 5652 AX Eindhoven, The Netherlands

Tel. +31 630 070 625, Fax: +31 402 920 069

E-mail: info@nordiclabs.nl

www.nordiclabs.nl