

<b>Description</b>	<b>IgG fraction of polyclonal goat antiserum to free and bound human Ig kappa light chain</b>	
<b>Product code</b>	GAHu/BJK(SD+HD)/7S	
<b>Biological origin</b>	Goat	
<b>Physical form</b>	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (pH 7.2).	
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity	
<b>Immunogen</b>	A pool of purified Bence Jones kappa proteins isolated from human urine. Freund's complete adjuvant is used in the first step of the immunization procedure.	
<b>Adsorption</b>	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other human serum proteins.	
<b>Purification</b>	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography.	
<b>Identity &amp; Specificity</b>	The reactivity of the antiserum is directed to the surface and hidden determinants of Ig kappa light chain. In immunoelectrophoresis this antiserum is reacting with polyclonal and monoclonal immunoglobulins of the kappa type, Bence Jones proteins as well as free light chains of the kappa type. This antiserum does not react with any other protein of human serum or plasma.	
<b>Cross-reactivity</b>	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins and their fragments, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
<b>Physicochemical characteristic</b>	IgG protein concentration is 10 mg/ml. No foreign proteins added.	
<b>Antibody titre</b>	Precipitin titre not less than 1:64 when tested against normal human serum in agar immunodiffusion block titration.	
<b>Intended uses</b>	<p>This IgG preparation is intended for use in precipitating (gel diffusion) and non-precipitating antibody-binding immunoassays (indirect immunofluorescence, enzyme immunoassays for the detection of polyclonal immunoglobulins, purified monoclonal immunoglobulins of the kappa type as well as free light chains or Bence Jones proteins of the kappa type in serum or other body fluids.; to prepare an insoluble immunoaffinity adsorbent or a solid phase catching antibody reagent by coupling to an artificial carrier; for labeling with a marker of the user's own choice.</p> <p><i>When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration.</i></p> <p>Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays between 1:500 and 1:5,000.</p>	
<b>Handling</b>	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.	
<b>Packing</b>	Vial with 10 mg lyophilized IgG (7S) preparation.	
<b>Storage / shelf life</b>	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
<b>Caution</b>	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

**NORDIC IMMUNOLOGICAL LABORATORIES**  
 Langendijk 25, 5652 AX Eindhoven, The Netherlands  
 Tel. +31 630 070 625, Fax: +31 402 920 069  
 E-mail: info@nordiclabs.nl  
 www.nordiclabs.nl