

Description	Precipitating polyclonal goat antiserum to free human Ig lambda chain	
Product code	GAHu/BJL(HD)	
Biological origin	Goat	
Physical form	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (pH 7.2).	
Preservative	No preservative added, as it may interfere with the antibody activity	
Immunogen	A pool of purified Bence Jones lambda proteins isolated from human urine. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with human serum proteins. Further adsorption by stepwise soluble adsorption to eliminate antibodies reacting with bound lambda chain.	
Identity & Specificity	The reactivity of the antiserum is directed to the hidden determinants of Ig lambda light chain. In immunoelectrophoresis this antiserum is only reacting with Bence Jones proteins as well as free light chains of the lambda type. No reaction is obtained with intact polyclonal or monoclonal immunoglobulins.	
Cross-reactivity	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins and their fragments, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
Physicochemical characteristic	Goat IgG protein concentration is about 25 mg/ml. No foreign proteins added.	
Intended use	To detect and identify free light chain type and Bence Jones protein in serum or urine by immunoelectrophoresis or double radial immunodiffusion. It is of great importance as an aid in the diagnosis of B cell malignancies such as multiple myeloma and non-Hodgkin lymphoma, and in monitoring their therapy. In immunoelectrophoresis use 1.2 µl of sample in a well with a diameter of 1 mm against 60 µl antiserum in a 20 mm long channel. Electrophoresis for 1 hour, 20 mA. The detection limit of this technique in serum is about 25 mg/l; in the case 100 times concentrated urine is used the detection limit can be brought down to 0.5 mg/l. The fact that in about 40% of the case the Bence Jones protein was only detectable in either serum or urine emphasizes the necessity to investigate both fluids.	
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles. Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.	
Packing	Vial with 1 ml lyophilized antiserum.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Caution	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl