

| | | | | | | | | | | | | | | | | | |
|-----------------------------------|---|-------------------|----------|----------|---------|----------|---------|-------|--------------|-----------|----------|---------|-----------|-----------|---------|-------|----------|
| Description | Precipitating polyclonal goat antiserum to human IgG, Fc specific | | | | | | | | | | | | | | | | |
| Product code | GAHu/IgG(Fc) | | | | | | | | | | | | | | | | |
| Biological origin | Goat | | | | | | | | | | | | | | | | |
| Physical form | Delipidated, heat inactivated, lyophilized, stable whole antiserum | | | | | | | | | | | | | | | | |
| Preservative | No preservative added. | | | | | | | | | | | | | | | | |
| Immunogen | Highly purified Fc fragment of normal IgG isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. | | | | | | | | | | | | | | | | |
| Adsorption | Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum. | | | | | | | | | | | | | | | | |
| Identity & Specificity | The reactivity of the antiserum is restricted to the Fc part of IgG. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal human serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgG. No precipitation reaction is obtained with purified IgA, IgM, and IgG/Fab fragments. | | | | | | | | | | | | | | | | |
| Cross-reactivity | <p>Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic determinants. The degree of cross-reactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiserum has been tested for cross-reactivity in double radial immunodiffusion with the following results:</p> <table border="0" style="margin-left: 40px;"> <tr> <td>bovine -</td> <td>dog -</td> <td>horse -</td> <td>pigeon -</td> <td>sheep -</td> </tr> <tr> <td>cat -</td> <td>guinea pig -</td> <td>monkey ++</td> <td>rabbit -</td> <td>swine -</td> </tr> <tr> <td>chicken -</td> <td>hamster -</td> <td>mouse -</td> <td>rat -</td> <td>turkey -</td> </tr> </table> <p>Three monkey sera were tested. A reaction with Rhesus monkey, Cynomolgus and Baboon have been found.</p> | | bovine - | dog - | horse - | pigeon - | sheep - | cat - | guinea pig - | monkey ++ | rabbit - | swine - | chicken - | hamster - | mouse - | rat - | turkey - |
| bovine - | dog - | horse - | pigeon - | sheep - | | | | | | | | | | | | | |
| cat - | guinea pig - | monkey ++ | rabbit - | swine - | | | | | | | | | | | | | |
| chicken - | hamster - | mouse - | rat - | turkey - | | | | | | | | | | | | | |
| Protein concentration | Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. | | | | | | | | | | | | | | | | |
| Antibody titre | Precipitin titre 1:64 when tested against pooled normal human serum in agar-block immunodiffusion titration. | | | | | | | | | | | | | | | | |
| Intended use | In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgG in human serum or other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of human IgG from serum or plasma. | | | | | | | | | | | | | | | | |
| Directions for use | In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted in 2 mm diameter peripheral wells). | | | | | | | | | | | | | | | | |
| Packing | Vial with 1 ml lyophilized antiserum. | | | | | | | | | | | | | | | | |
| Storage / shelf life | Lyophilized at +4°C | at least 10 years | | | | | | | | | | | | | | | |
| | reconstituted at or below -20°C | 3-5 years | | | | | | | | | | | | | | | |
| | reconstituted at +4°C | 7 days | | | | | | | | | | | | | | | |
| Handling | The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. | | | | | | | | | | | | | | | | |
| Caution | This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only. | | | | | | | | | | | | | | | | |

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl