

<b>Description</b>	<b>Purified IgG fraction of polyclonal goat antiserum to mouse fibrinogen</b>
<b>Product code</b>	GAM/Fbg/7S
<b>Biological origin</b>	Goat
<b>Physical form</b>	Purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2)
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity.
<b>Immunogen</b>	Fibrinogen (clotting factor I) is a heat labile beta glycoprotein present in plasma. It is the precursor of fibrin, which is the key protein constituting the network of the blood clot. Thrombin converts fibrinogen to fibrin by limited proteolysis. Fibrin monomers polymerize to fibrin which is stabilized by cross-linking. Fibrinogen is isolated from fresh plasma after removing prothrombin. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Purification</b>	The IgG (7S) fraction is isolated and purified from the antiserum and contains the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis and double radial immunodiffusion.
<b>Adsorption</b>	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
<b>Identity &amp; Specificity</b>	The reactivity of the antiserum is restricted to fibrinogen. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal mouse plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified fibrinogen. No reaction is obtained with any other plasma protein component or serum. However, the antiserum may also react with fibrin monomers, circulating fibrinopeptides and fibrin degradation products.
<b>Cross-reactivity</b>	The antiserum does not cross-react with any other component of mouse plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.
<b>Physicochemical characteristic</b>	IgG protein concentration 9.8 mg/ml. No foreign proteins added.
<b>Intended use</b>	As unlabelled primary or secondary antibody reagent for the indirect detection of fibrinogen in mouse cells, tissues and body fluids in immunofluorescence and immunoenzyme methods; for the production of immunoconjugates with a selected marker; to prepare insoluble immunoaffinity adsorbents by coupling to an artificial carrier; as catching or detection reagent in non-isotopic methodology and solid phase immunochemistry (e.g. ELISA, Western blotting). <i>When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration.</i> Typical working dilutions in histochemistry are usually between 1:50 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays between 1:500 and 1:5,000.
<b>Directions for use</b>	The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.
<b>Packing</b>	Vial with 9.8 mg lyophilized IgG (7S) fraction.
<b>Storage / shelf life</b>	Lyophilized at +4°C at least 10 years reconstituted at or below -20°C 3-5 years reconstituted at +4°C 7 days
<b>Caution</b>	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.

**NORDIC IMMUNOLOGICAL LABORATORIES**

Langendijk 25, 5652 AX Eindhoven, The Netherlands

Tel. +31 630 070 625, Fax: +31 402 920 069

E-mail: [info@nordiclabs.nl](mailto:info@nordiclabs.nl)[www.nordiclabs.nl](http://www.nordiclabs.nl)