

Description	Purified IgG fraction of polyclonal goat antiserum to mouse immunoglobulins IgG, IgA, IgM, heavy and light chains																
Product code	GAM/Ig/7S																
Biological origin	Goat																
Physical form	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).																
Preservative	No preservative added, as it may interfere with the antibody activity.																
Immunogen	Highly purified IgG, including all subclasses, and pools of homogenous IgA and IgM isolated from mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.																
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG (7S) fraction by DEAE-chromatography.																
Adsorption	Immunoaffinity adsorbed using insolubilized Ig-depleted mouse serum as required, to eliminate antibodies reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.																
Identity & Specificity	The defined antibody specificity is directed to the major isotypes of the mouse immunoglobulin system (classes, subclasses and L chain types) including antibody to common determinants, to class and subclass-specific determinants, and to the surface determinants of the common Fab portion as tested against as tested in immunoelectrophoresis and double radial immunodiffusion (Ouchterlony).																
Cross-reactivity	<p>Inter-species cross-reactivity is a normal feature of antibodies to mammalian immunoglobulins, since homologous proteins of different species frequently share antigenic determinants. The degree of cross-reactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiserum fraction has been tested for cross-reactivity by double radial immunodiffusion against several species sera with the following results:</p> <table border="0" style="margin-left: 40px;"> <tr> <td>bovine -</td> <td>duck -</td> <td>human -</td> <td>rat +</td> </tr> <tr> <td>cat -</td> <td>guinea pig +</td> <td>monkey -</td> <td>sheep -</td> </tr> <tr> <td>chicken -</td> <td>hamster ±</td> <td>pigeon -</td> <td>swine -</td> </tr> <tr> <td>dog -</td> <td>horse -</td> <td>rabbit -</td> <td>turkey -</td> </tr> </table>	bovine -	duck -	human -	rat +	cat -	guinea pig +	monkey -	sheep -	chicken -	hamster ±	pigeon -	swine -	dog -	horse -	rabbit -	turkey -
bovine -	duck -	human -	rat +														
cat -	guinea pig +	monkey -	sheep -														
chicken -	hamster ±	pigeon -	swine -														
dog -	horse -	rabbit -	turkey -														
Physicochemical characteristic	IgG protein concentration 10 mg/ml. No foreign proteins added.																
Intended use	<p>The cytochemical grade of GAM/Ig/7S allows its use in different types of highly sensitive immunoassays on appropriately treated cell and tissue substrates; in radioimmunoassay; for the production of immunoconjugates with a selected marker; to prepare immunoaffinity adsorbents by coupling to an artificial carrier; in non-isotopic methodology based on solid phase immunochemistry (e.g. ELISA), both as catching antibody and detection reagent; in Western blotting.</p> <p><i>This product is not pre-diluted. The optimum working dilution of each product should be established by titration before being used.</i></p> <p>Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays are between 1:500 and 1:5,000.</p>																
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.																
Packing	Vial with 10 mg lyophilized IgG(7S) fraction.																
Storage / shelf life	<table border="0" style="margin-left: 40px;"> <tr> <td>Lyophilized at +4°C</td> <td>at least 10 years</td> </tr> <tr> <td>reconstituted at or below -20°C</td> <td>3-5 years</td> </tr> <tr> <td>reconstituted at +4°C</td> <td>7 days</td> </tr> </table>	Lyophilized at +4°C	at least 10 years	reconstituted at or below -20°C	3-5 years	reconstituted at +4°C	7 days										
Lyophilized at +4°C	at least 10 years																
reconstituted at or below -20°C	3-5 years																
reconstituted at +4°C	7 days																
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.																

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl