

Description	Biotin-conjugated IgG fraction of polyclonal goat antiserum to mouse IgG2a, subclass specific	
Product code	GAM/IgG2a/Bio	
Biological origin	Goat	
Physical form	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).	
Preservative	No preservative added, as it may interfere with the antibody activity.	
Immunogen	Pools of purified homogenous IgG2a isolated from mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation and purification of the IgG (7S) fraction containing the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Identity & Specificity	The reactivity of the antiserum is directed to the subclass IgG2a. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in mouse serum, as tested by immunoelectrophoresis and double radial immunodiffusion.	
Cross-reactivity	This immunoconjugate is not species-specific since inter-species cross-reactivity is a normal feature of antisera to immunoglobulins. However this conjugate has been passed over appropriate immuno-adsorbents to remove antibodies cross-reacting with human immunoglobulins. This renders it specific for use in test systems containing material of human origin (e.g. human tissue/mouse monoclonal antibody to a human tissue constituent/anti mouse Ig isotype-specific immunoconjugate).	
Physicochemical characteristics	IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio (B/P) is approximately 6.9. No foreign proteins added.	
Marker	N-Hydroxysuccinimidobiotin.	
Conjugation procedure	A proprietary technique for the binding of biotin, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life.	
Intended use	<p>In immunocytochemical and immunohistochemical staining for the detection of IgG2a at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating IgG2a antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of mouse origin known to be of the IgG2a isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgG2a in mouse serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used</p> <p><i>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</i></p> <p>Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays between 1:1.000 and 1:8.000 depending on the method used.</p>	
Handling	<p>The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.</p>	
Packing	Vial with 1 ml lyophilized immunoconjugate.	
Storage / shelf life	Lyophilized at +4°C reconstituted at or below -20°C reconstituted at +4°C	at least 10 years 3-5 years 7 days
Caution	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl