Description	**Purified IgG fraction of polyclonal goat antiserum to monkey immunoglobulins IgG, IgA, IgM, heavy and light chains**
Product code	GAMon/Ig/7S
Biological origin	Goat
Physical form	Purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).
Preservative	No preservative added, as it may interfere with the antibody activity.
Immunogen	Highly purified IgG and pools of homogenous IgA and IgM isolated from monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG (7S) fraction by DEAE-chromatography.
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
Identity & Specificity	The reactivity of the antiserum is directed to the major isotypes of the monkey immunoglobulin system (classes and both light chain types) including antibodies to common determinants, to class and to the surface determinants of the common Fab portion, as tested by immunoelectrophoresis and double radial immunodiffusion against pooled serum and purified immunoglobulins. In immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations against normal monkey plasma and serum, the characteristic IgG, IgA and IgM precipitin lines are obtained.
Cross-reactivity	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Precipitation reactions have been observed with immunoglobulins in serum of other old-world monkeys, including Cercopithecus, Cynomolgus and Baboon. The product may also react with other species as has been observed for chimpanzee and man.
Physicochemical characteristic	IgG protein concentration 10 mg/ml. No foreign proteins added.
Intended use	This IgG fraction allows the use in different types of highly sensitive immunoassays on appropriately treated cell and tissue substrates; in radioimmunoassay; for the production of immunoconjugates with a selected marker; to prepare immunoaffinity adsorbents by coupling to an artificial carrier; in non-isotopic methodology based on solid phase immunochemistry (e.g. ELISA), both as catching antibody and detection reagent; in Western blotting. *This product is not pre-diluted. The optimum working dilution of each product should be established by titration before being used.* Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays are between 1:1,000 and 1:5,000.
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.
Packing	Vial with 10 mg lyophilized IgG (7S) fraction.
Storage / shelf life	Lyophilized at +4°C — at least 10 years reconstituted at or below -20°C — 3-5 years reconstituted at +4°C — 7 days
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For *in vitro* laboratory research purposes only.

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl

© Nordic Immunological Laboratories 2004