

<b>Description</b>	<b>Purified IgG fraction of polyclonal goat antiserum to monkey lactoferrin</b>						
<b>Product code</b>	GAMon/Lfr/7S						
<b>Biological origin</b>	Goat						
<b>Physical form</b>	Purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2)						
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity.						
<b>Immunogen</b>	<p>Exocrine organs produce various secretions, each with its characteristic function. Proteins found in secretions may be divided into two groups: those specific for the particular secretion, and plasma proteins independent of the type of exocrine cells. Lactoferrin belongs to the first group. It is an iron containing protein with a molecular weight of 75,000 and it is antigenically different from transferrin. Lactoferrin has a slight anti-microbial action. Originally identified in milk, its presence has also been demonstrated in other secretions as saliva, semen and tears. The immunogen has been isolated from rhesus monkey milk.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>						
<b>Adsorption</b>	Immunoaffinity adsorbed using insolubilized fractions of monkey serum and lactoferrin-depleted monkey milk as required, to eliminate antibodies reacting with other monkey serum or milk proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.						
<b>Purification</b>	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography.						
<b>Identity &amp; Specificity</b>	In immunoelectrophoresis against monkey milk a single precipitin line is obtained. The antiserum does not react with any other protein component of monkey serum or plasma.						
<b>Cross-reactivity</b>	The antiserum does not cross-react with any other monkey plasma proteins as tested in gel-diffusion techniques. Inter-species cross-reactivity is a normal feature of antibodies to plasma or milk proteins, since homologous proteins of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail, however a strong cross reaction with lactoferrin in human milk has been observed.						
<b>Physicochemical characteristics</b>	IgG protein concentration 10 mg/ml. No foreign proteins added.						
<b>Intended use</b>	<p>In enzyme-immunocytochemical and immunohistochemical techniques for the indirect detection of monkey lactoferrin at the cellular and subcellular level in appropriately treated cell and tissue substrates; as catching reagent in non-isotopic methodology and solid phase immunochemistry (e.g. ELISA, Western blotting).</p> <p><i>The optimum working dilution of the product should be established by titration before being used.</i></p> <p>Working dilutions for histochemical and cytochemical use are usually between 1:50 and 1:500.</p>						
<b>Directions for use</b>	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.						
<b>Packing</b>	Vial with 10 mg lyophilized IgG (7S) fraction.						
<b>Storage / shelf life</b>	<table><tr><td>Lyophilized at +4°C</td><td>at least 10 years</td></tr><tr><td>reconstituted at or below -20°C</td><td>3-5 years</td></tr><tr><td>reconstituted at +4°C</td><td>7 days</td></tr></table>	Lyophilized at +4°C	at least 10 years	reconstituted at or below -20°C	3-5 years	reconstituted at +4°C	7 days
Lyophilized at +4°C	at least 10 years						
reconstituted at or below -20°C	3-5 years						
reconstituted at +4°C	7 days						
<b>Caution</b>	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.						

**NORDIC IMMUNOLOGICAL LABORATORIES**

Langendijk 25, 5652 AX Eindhoven, The Netherlands

Tel. +31 630 070 625, Fax: +31 402 920 069

E-mail: [info@nordiclabs.nl](mailto:info@nordiclabs.nl)[www.nordiclabs.nl](http://www.nordiclabs.nl)