

<b>Description</b>	<b>Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal goat antiserum to C3c fragment of rabbit complement factor C3</b>	
<b>Product code</b>	GAR/C3c/FITC	
<b>Biological origin</b>	Goat	
<b>Physical form</b>	Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2)	
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity.	
<b>Immunogen</b>	C3c is the major fragment resulting from C3 cleavage by C3 convertase and factor I. It is composed of an intact beta chain bound to two fragments of the alpha chain. C3c is isolated and purified from pooled normal rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure.	
<b>Purification</b>	The IgG (7S) fraction is isolated and purified from the antiserum and contains the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis and double radial immunodiffusion.	
<b>Adsorption</b>	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
<b>Identity &amp; Specificity</b>	In immunoelectrophoresis against fresh rabbit serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the alpha-2 region, demonstrating a gradient. In old serum containing totally activated C3 a single precipitin line in the alpha-2 region is obtained. Antisera to C3c cab also react with the fragments C3b, C3bi and smaller fragments, since they all carry antigenic determinants of the C3c domain. The product does not react with any other proteins component of rabbit serum or plasma.	
<b>Cross-reactivity</b>	The antiserum does not cross-react with any other component of rabbit plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
<b>Physicochemical characteristics</b>	IgG protein concentration 15.5 mg/ml. Fluorescein/IgG protein molar ratio (F/P) approximately 1.2. No foreign proteins added.	
<b>Fluorescent marker</b>	Fluorescein isothiocyanate isomer 1 (FITC). Excitation: 492 nm, emission: 515 nm.	
<b>Intended use</b>	The fluorescent immunoconjugate to rabbit C3c is used to determine the presence and pattern of C3 in tissue lesions using immunohistochemical staining techniques. Locally deposited immune complexes in tissue usually contain complement, pointing to activation of the classical pathway. Complement activation in vivo implies active disease and may contribute to the elicitation of the pathogenesis and he extent of tissue destruction. Sometimes the diagnosis can be based on directly on laboratory findings. <i>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</i> Working dilutions are usually between 1:20 and 1:80.	
<b>Handling</b>	The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.	
<b>Packing</b>	Vial with 1 ml lyophilized immunoconjugate.	
<b>Storage / shelf life</b>	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
<b>Caution</b>	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.	

### NORDIC IMMUNOLOGICAL LABORATORIES

Langendijk 25, 5652 AX Eindhoven, The Netherlands  
 Tel. +31 630 070 625, Fax: +31 402 920 069  
 E-mail: info@nordiclabs.nl  
 www.nordiclabs.nl