Description	**Biotin-conjugated purified monoclonal mouse antibody to human IgD, isotype specific**
Product code	MAHu/IgD/Bio
Biological origin	Mouse, clone NI 158
Mouse isotype	IgG1κ
Physical form	Purified monoclonal mouse IgG1κ conjugated with biotin, lyophilized from a solution in phosphate buffered saline (pH7.2).
Preservative	No preservative added, as it may interfere with the antibody activity. No foreign protein added.
Immunogen	Highly purified polyclonal IgD isolated from human serum.
Identity & Specificity	The reactivity of the antiserum is restricted to an isotype specific determinant of the IgD molecule as tested in direct binding enzyme immunoassay, immunoblotting, immunoprecipitation and direct immunoperoxidase staining of cytoplasmic Ig.
Cross-reactivity	The antiserum does not react with any other component of the human Ig system or any other human plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species.
Physicochemical characteristics	IgG concentration is 0.4 mg/ml. Biotin/IgG protein molar ratio (B/P) approximately 5.0. No foreign proteins added.
Marker	N-Hydroxysuccinimidobiotin.
Conjugation procedure	Conjugation is carried out using a proprietary technique for the coupling of biotin, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. No foreign protein has been added. The conjugate is lyophilized to assure stability and long shelf life.
Intended use	To identify the presence of IgD in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, immunoperoxidase staining of cytoplasmic IgD, and immunoblotting. As a second step an avidin or streptavidin conjugate of the customer's choice have to be used. The optimum working dilution is an assay-related characteristic. It may vary widely and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:50 to 1:200; in ELISA from 1:500 upwards; in Western blotting from 1:1,000 upwards. These data should be interpreted as general recommendations only. Working dilutions may vary widely, strongly depending on the test conditions.
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4ºC, not refrozen, and preferably used the same day.
Packing	Vial with 0.5 ml lyophilized immunoconjugate.
Storage / shelf life	Lyophilized at +4ºC — at least 10 years reconstituted at or below -20°C — 3-5 years reconstituted at +4°C — 7 days
Caution	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For *in vitro* research purposes only.

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl

© Nordic Immunological Laboratories 2004