Description	**Fluorescein isothiocyanate-conjugated purified monoclonal mouse antibody to human IgG1, subclass specific**
Product code	MAHu/IgG1/FITC
Biological origin	Mouse, clone NI 132 (HP 6186)
Mouse isotype	IgG1κ
Physical form	Purified monoclonal mouse IgG1κ conjugated with FITC, lyophilized from a solution in phosphate buffered saline (pH7.2).
Preservative	No preservative added, as it may interfere with the antibody activity. No foreign protein added.
Immunogen	Highly purified monoclonal IgG1 isolated from human serum.
Identity & Specificity	The reactivity of the antiserum is restricted to a subclass specific determinant on the Fc part of the IgG1 molecule as tested in direct binding enzyme immunoassay, immunoblotting, immuno-precipitation and direct immunoperoxidase staining of cytoplasmic Ig.
Cross-reactivity	The antiserum does not react with any other component of the human Ig system or any other human plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species.
Physicochemical characteristics	IgG concentration is 0.4 mg/ml. Fluorochrome/IgG protein molar ratio (F/P) approximately 2.7. No foreign proteins added.
Fluorescent marker	Fluorescein isothiocyanate isomer 1. Excitation: 492 nm, emission: 515 nm.
Conjugation procedure	A proprietary technique for the binding to FITC is used, followed by several purification steps to remove free reactants and protein aggregates. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life.
Intended use	To identify the presence of IgG1 in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, indirect immunoperoxidase staining of cytoplasmic IgG1, and immunoblotting using a peroxidase labelled monoclonal antibody against FITC. The optimum working dilution is an assay-related characteristic. It may vary widely and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:10 to 1:50; in ELISA from 1:200 upwards; in Western blotting from 1:400 upwards. Working dilutions may vary widely, strongly depending on the test conditions. These data should be interpreted as general recommendations only.
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4ºC, not refrozen, and preferably used the same day.
Packing	Vial with 0.5 ml lyophilized immunoconjugate.
Storage / shelf life	Lyophilized at +4ºC — at least 10 years reconstituted at or below -20°C — 3-5 years reconstituted at +4°C — 7 days
Caution	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For *in vitro* research purposes only.
Reference	Evaluation of monoclonal antibodies having specificity for human IgG subclasses: Results of an IUIS/WHO collaborative study. Immunology Letters **10** (1985), 223-252. Jefferis R, Reimer C B, Skavril F, de Lange G, Ling N R, Lowe L, Walker M R, Philips D J, Aloisio C H, Wells T W, Vaerman J, Magnusson C G, Kubagawa H, Cooper M, Vartdal F, Vandvik B, Haaijman J J, Makela O, Sarnesto A, Lando Z, Gergely J, Rainavölgyi E, László G, Radl J and Molinaro G A.

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl

© Nordic Immunological Laboratories 2004