

<b>Description</b>	<b>Monoclonal mouse antiserum to human IgG, isotype specific</b>	
<b>Product code</b>	MAHu/IgGc	
<b>Biological origin</b>	Mouse, clones NI 335 and NI 343	
<b>Mouse isotype</b>	IgG1 $\kappa$	
<b>Physical form</b>	Delipidated, heat inactivated, lyophilized, stable whole ascites	
<b>Immunogen</b>	Highly purified IgG isolated from pooled human serum	
<b>Identity &amp; Specificity</b>	The reactivity of the antiserum is restricted to isotype specific determinants on the Fc part of the IgG molecule. The antiserum reacts with the 4 subclasses of IgG, as tested in indirect binding enzyme immunoassay, immunoblotting and indirect immunoperoxidase staining of cytoplasmic Ig.	
<b>Cross-reactivity</b>	The antiserum does not react with any other component of the human Ig system or any other human plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species.	
<b>Protein concentration</b>	IgG concentration is 1.0 mg/ml. No foreign proteins added.	
<b>Intended use</b>	To identify the presence of IgG in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as radio immuno assay, ELISA, indirect immunoperoxidase and immunofluorescence staining and immunoblotting of cytoplasmic IgG. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:50 to 1:200; in ELISA from 1:500 upwards; in Western blotting from 1:1,000 upwards. These data should be interpreted as general recommendations only.	
<b>Handling</b>	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized ascites by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted ascites should be stored at +4°C, not refrozen, and preferably used the same day.	
<b>Packing</b>	Vial with 0.5 ml lyophilized ascites.	
<b>Storage / shelf life</b>	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
<b>Caution</b>	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.	

**NORDIC IMMUNOLOGICAL LABORATORIES**

Langendijk 25, 5652 AX Eindhoven, The Netherlands  
Tel. +31 630 070 625, Fax: +31 402 920 069  
E-mail: info@nordiclabs.nl  
www.nordiclabs.nl