Description	**Monoclonal mouse antibody to human IgG, isotype specific**
Product code	MAHu/IgGc/MAb
Biological origin	Mouse, clones NI 335 and NI 343
Mouse isotype	IgG1κ
Physical form	Purified mouse IgG1κ lyophilized from a solution in phosphate buffered saline (pH 7.2).
Preservative	No preservative added, as it may interfere with the antibody activity. No foreign protein added.
Immunogen	Highly purified monoclonal IgG isolated from human serum
Identity & Specificity	The reactivity of the antiserum is restricted to isotype specific determinants on the Fc part of the IgG molecule. It reacts with the 4 subclasses of human IgG, as tested in indirect binding enzyme immunoassay, immunoblotting, immunoprecipitation and indirect immunoperoxidase staining of cytoplasmic Ig.
Cross-reactivity	The antiserum does not react with any other component of the human Ig system or any other human plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species.
Protein concentration	IgG concentration is 0.4 mg/ml. No foreign proteins added.
Intended use	To identify the presence of IgG in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, indirect immunoperoxidase and indirect immunofluorescence staining of cytoplasmic IgG, and immunoblotting. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:50 to 1:200; in ELISA from 1:500 upwards; in Western blotting from 1:1,000 upwards. These data should be interpreted as general recommendations only.
Handling	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4ºC, not refrozen, and preferably used the same day.
Packing	Vial with 0.5 ml lyophilized monoclonal antibody.
Storage / shelf life	Lyophilized at +4ºC — at least 10 years reconstituted at or below -20°C — 3-5 years reconstituted at +4°C — 7 days
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For *in vitro* research purposes only.
Reference	Evaluation of monoclonal antibodies having specificity for human IgG subclasses: Results of the 2nd IUIS/WHO collaborative study. Immunology Letters **31** (1992), 143-168. Jefferis R, Reimer C B, Skavril F, de Lange G, Goodall D M, Bentley T L, Phillips D J, Vlug A, Harada S, Radl J , Claassen E, Boersma J A, and Coolen J.

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl

© Nordic Immunological Laboratories 2006