

| | | |
|----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|
| Description | Peroxidase conjugated purified monoclonal mouse antibody to human lactoferrin | |
| Product code | MAHu/Lfr/PO | |
| Biological origin | Mouse, clone NI 25 | |
| Mouse isotype | IgG1 κ | |
| Physical form | Purified monoclonal mouse IgG1 κ conjugated with horseradish peroxidase, lyophilized from a solution in phosphate buffered saline (pH7.2). | |
| Preservative | No preservative added, as it may interfere with the antibody activity. No foreign protein added. | |
| Immunogen | Highly purified lactoferrin isolated from human milk. | |
| Identity & Specificity | The reactivity of the antiserum is restricted to lactoferrin as tested in indirect binding immunoblotting and indirect immunoperoxidase staining. | |
| Cross-reactivity | The antiserum does not react with any other component of the human milk or plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species. | |
| Physicochemical characteristics | IgG concentration is 0.4 mg/ml. Peroxidase/IgG protein molar ratio (E/P) approximately 1.7. No foreign proteins added. | |
| Enzyme marker | Horseradish peroxidase enriched for isoenzyme C (RZ=3.2). | |
| Conjugation procedure | Conjugation is carried out using a proprietary modification of the periodate method, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. No foreign protein has been added. The conjugate is lyophilized to assure stability and long shelf life. | |
| Intended use | To identify the presence of lactoferrin in human milk, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, indirect immunoperoxidase and indirect immunofluorescence staining and immunoblotting. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:20 upwards; in ELISA from 1:100 upwards; in Western blotting from 1:250 upwards. These data should be interpreted as general recommendations only. | |
| Handling | The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4°C, not refrozen, and preferably used the same day. | |
| Packing | Vial with 0.5 ml lyophilized immunoconjugate. | |
| Storage / shelf life | Lyophilized at +4°C | at least 10 years |
| | reconstituted at or below -20°C | 3-5 years |
| | reconstituted at +4°C | 7 days |
| Caution | This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only. | |

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl