

<b>Description</b>	<b>Biotin-conjugated purified monoclonal mouse antibody to human secretory component, free or bound</b>
<b>Product code</b>	MAHu/SC/Bio
<b>Biological origin</b>	Mouse, clone NI 194-4 (A89-39)
<b>Mouse isotype</b>	IgG1 $\kappa$
<b>Physical form</b>	Purified monoclonal mouse IgG1 $\kappa$ conjugated with biotin, lyophilized from a solution in phosphate buffered saline (pH7.2).
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity. No foreign protein added.
<b>Immunogen</b>	Highly purified secretory component isolated from human milk.
<b>Identity &amp; Specificity</b>	The reactivity of the antiserum is restricted to secretory component as tested in indirect binding immunoblotting and indirect immunoperoxidase staining.
<b>Cross-reactivity</b>	The antiserum does not react with any other component of the human milk or plasma protein as tested. This antiserum has not been tested for cross-reactivity with other species.
<b>Physicochemical characteristics</b>	IgG concentration is 0.4 mg/ml. Biotin/IgG protein molar ratio (B/P) approximately 1.4. No foreign proteins added.
<b>Marker</b>	N-Hydroxysuccinimidobiotin.
<b>Conjugation procedure</b>	Conjugation is carried out using a proprietary technique for the coupling of biotin, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. No foreign protein has been added. The conjugate is lyophilized to assure stability and long shelf life.
<b>Intended use</b>	To identify the presence of free or bound secretory component in human milk, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, indirect immunoperoxidase and indirect immunofluorescence staining and immunoblotting. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:20 upwards; in ELISA from 1:25 upwards; in Western blotting from 1:50 upwards. These data should be interpreted as general recommendations only.
<b>Handling</b>	The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4°C, not refrozen, and preferably used the same day.
<b>Packing</b>	Vial with 0.5 ml lyophilized immunoconjugate.
<b>Storage / shelf life</b>	Lyophilized at +4°C at least 10 years reconstituted at or below -20°C 3-5 years reconstituted at +4°C 7 days
<b>Caution</b>	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.

**NORDIC IMMUNOLOGICAL LABORATORIES**

Langendijk 25, 5652 AX Eindhoven, The Netherlands

Tel. +31 630 070 625, Fax: +31 402 920 069

E-mail: info@nordiclabs.nl

www.nordiclabs.nl