

Description	Precipitating polyclonal rabbit antiserum to human immunoglobulins, IgG, IgA, IgM and IgD, heavy and light chains	
Product code	RAHu/Ig(GAMD)	
Biological origin	Rabbit	
Physical form	Delipidated, heat inactivated, lyophilized stable whole serum	
Preservative	No preservative added, as it may interfere with the antibody activity. No foreign protein added.	
Immunogen	Purified normal IgG and fractions containing IgA, IgM and IgD isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibody activity with other serum protein. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Identity & Specificity	The reactivity of the antiserum is directed to the major isotypes of the human immunoglobulin system (classes and both light chain types) including antibodies to common determinants, to class and to the surface determinants of the common Fab portion, as tested by immunoelectrophoresis and double radial immunodiffusion against pooled serum and purified immunoglobulins.	
Cross-reactivity	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
Protein concentration	Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added.	
Antibody titre	Precipitin titre not less than 1:64 when tested against normal human serum in agar block titration.	
Intended use	In precipitating techniques as immunoelectrophoresis, single and double radial immunodiffusion (Mancini, Ouchterlony) to identify and measure immunoglobulins; as secondary antibody to precipitate the immunoglobulin in normal human serum.	
Directions for use	In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.	
Handling	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.	
Packing	Vial with 1 ml lyophilized antiserum.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Caution	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl