

Description	IgG fraction of polyclonal rabbit antiserum to human lysozyme	
Product code	RAHu/Lys/7S	
Biological origin	Rabbit	
Physical form	Purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2)	
Preservative	No preservative added, as it may interfere with the antibody activity.	
Immunogen	Lysozyme, a bacteriolytic enzyme, has a molecular weight of about 15,000. Its concentration in plasma is low; in milk the concentration is reported to be between 5 and 220 mg/ml. Also tears contain a considerable amount of lysozyme. Highly purified lysozyme is isolated from pooled milk. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other with other plasma proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography.	
Identity & Specificity	The reactivity of the antiserum is restricted to lysozyme. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal human milk and tears a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified lysozyme. No reaction is obtained with any other plasma protein component or serum.	
Cross-reactivity	The antiserum does not cross-react with any other component of human plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
Physicochemical characteristics	IgG protein concentration 10 mg/ml. No foreign proteins added.	
Intended use	<p>The cytochemical grade allows the use in different types of highly sensitive immunoassays on appropriately treated cell and tissue substrates; in radioimmunoassay; for the production of immunoconjugates with a selected marker; to prepare immunoaffinity adsorbents by coupling to an artificial carrier; in non-isotopic methodology based on solid phase immunochemistry (e.g. ELISA), both as catching antibody and detection reagent; in Western blotting.</p> <p>This product is not pre-diluted. The optimum working dilution of each product should be established by titration before being used.</p> <p>Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays are between 1:500 and 1:5,000.</p>	
Directions for use	The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.	
Packing	Vial with 1 ml lyophilized IgG fraction.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl