

| | | |
|-----------------------------------|---|-------------------|
| Description | Precipitating polyclonal rabbit antiserum to monkey IgG, Fc specific | |
| Product code | RAMon/IgG(Fc) | |
| Biological origin | Rabbit | |
| Physical form | Delipidated, heat inactivated, lyophilized, stable whole antiserum | |
| Preservative | No preservative added | |
| Immunogen | Highly purified normal IgG isolated from pooled rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure. | |
| Adsorption | Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum. | |
| Identity & Specificity | The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal monkey serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgG. No precipitation reaction is obtained with purified IgA, IgM, and IgG/Fab fragments. | |
| Cross-reactivity | The antiserum does not cross react with any other component of the monkey immunoglobulin system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Precipitation reactions have been observed with IgG in serum of other old-world monkeys, including Cercopithecus, Cynomolgus and baboon. The antiserum may also react with IgG of other species as has been observed for chimpanzee and man. | |
| Protein concentration | Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. | |
| Antibody titre | Precipitin titre 1:64 when tested against pooled normal monkey serum in agar-block immunodiffusion titration. | |
| Intended use | In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgG in monkey serum or other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of monkey IgG from serum or plasma. | |
| Directions for use | In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted in 2 mm diameter peripheral wells). | |
| Packing | Vial with 1 ml lyophilized antiserum. | |
| Storage / shelf life | Lyophilized at +4°C | at least 10 years |
| | reconstituted at or below -20°C | 3-5 years |
| | reconstituted at +4°C | 7 days |
| Handling | The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. | |
| Caution | This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only. | |

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl