

<b>Description</b>	<b>Precipitating polyclonal rabbit antiserum to monkey IgG, heavy and light chains</b>																
<b>Product code</b>	RAMon/IgG(H+L)																
<b>Biological origin</b>	Rabbit																
<b>Physical form</b>	Delipidated, heat inactivated lyophilized stable whole serum																
<b>Preservative</b>	No preservative added, as it may interfere with the antibody activity. No foreign protein added.																
<b>Immunogen</b>	Purified normal IgG isolated from pooled rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.																
<b>Adsorption</b>	No adsorption required.																
<b>Identity &amp; Specificity</b>	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in monkey serum, as tested by immunoelectrophoresis and double radial immunodiffusion.																
<b>Cross-reactivity</b>	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Precipitation reactions have been observed with IgG in serum of other old-world monkeys, including Cercopithecus, Cynomolgus and Baboon. The conjugate may also react with IgG of other species as has been observed for chimpanzee and man. Cross-reactivity of this antiserum has been tested in double radial immunodiffusion with the following results: <table border="0" style="margin-left: 20px;"> <tr> <td>bovine +</td> <td>dog +</td> <td>hamster +</td> <td>mouse +</td> <td>sheep +</td> </tr> <tr> <td>cat +</td> <td>goat +</td> <td>horse +</td> <td>pigeon -</td> <td>swine +</td> </tr> <tr> <td>chicken -</td> <td>guinea pig +</td> <td>human ++</td> <td>rat +</td> <td>turkey -</td> </tr> </table>		bovine +	dog +	hamster +	mouse +	sheep +	cat +	goat +	horse +	pigeon -	swine +	chicken -	guinea pig +	human ++	rat +	turkey -
bovine +	dog +	hamster +	mouse +	sheep +													
cat +	goat +	horse +	pigeon -	swine +													
chicken -	guinea pig +	human ++	rat +	turkey -													
<b>Protein concentration</b>	Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added.																
<b>Antibody titre</b>	Precipitin titre not less than 1:64 when tested against normal monkey serum in agar block titration.																
<b>Intended use</b>	In precipitating techniques as immunoelectrophoresis and double radial immunodiffusion to identify the presence of IgG; as secondary antibody to precipitate the immunoglobulin in normal monkey serum; to prepare an immunoadsorbent for the production of Ig-free monkey serum, plasma or non-Ig preparations.																
<b>Directions for use</b>	In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.																
<b>Handling</b>	The lyophilised antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.																
<b>Packing</b>	Vial with 1 ml lyophilized antiserum.																
<b>Storage / shelf life</b>	Lyophilised at +4°C	at least 10 years															
	reconstituted at or below -20°C	3-5 years															
	reconstituted at +4°C	7 days															
<b>Caution</b>	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.																

**NORDIC IMMUNOLOGICAL LABORATORIES**  
 Langendijk 25, 5652 AX Eindhoven, The Netherlands  
 Tel. +31 630 070 625, Fax: +31 402 920 069  
 E-mail: info@nordiclabs.nl  
 www.nordiclabs.nl