

Description	IgG fraction of polyclonal rabbit antiserum to sheep IgM, Fc specific	
Product code	RASh/IgM(Fc)/7S	
Biological origin	Rabbit	
Physical form	Purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).	
Preservative	No preservative added, as it may interfere with the antibody activity.	
Immunogen	Purified normal IgM isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation by salt-precipitation and purification of the IgG fraction by DEAE-chromatography.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Identity & Specificity	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in sheep serum, as tested by immunoelectrophoresis and double radial immunodiffusion.	
Cross-reactivity	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail, however a strong reaction with goat IgM has been observed.	
Physicochemical characteristic	IgG protein concentration 10 mg/ml. No foreign proteins added.	
Antibody titre	Precipitin titre 1:64 when tested against pooled normal sheep serum in agar-block immunodiffusion titration.	
Intended use	<p>As unlabelled primary or secondary reagent for indirect detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to prepare conjugates of the user's own choice; to prepare an insoluble immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching antibody in non-isotopic methodology and solid phase immunochemistry.</p> <p><i>When applied in any cytochemical or histochemical staining procedure or solid phase coupling technique, the optimum concentration of the IgG preparation should be established by titration before being used.</i></p> <p>Typical working dilutions in histochemistry are usually between 1:50 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays between 1:500 and 1:5.000.</p>	
Handling	<p>The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.</p>	
Packing	Vial with 10 mg lyophilized IgG (7S) fraction.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Caution	This product should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl