

| | | | | | | | | | | | | | | | | | |
|-----------------------------------|--|-------------------|----------|----------|--------------|---------|-------|-------|--------|---------|----------|---------|-----------|--------|---------|----------|----------|
| Description | Precipitating polyclonal rabbit antiserum to swine IgG, heavy and light chains | | | | | | | | | | | | | | | | |
| Product code | RASw/IgG(H+L) | | | | | | | | | | | | | | | | |
| Biological origin | Rabbit | | | | | | | | | | | | | | | | |
| Physical form | Delipidated, heat inactivated lyophilized stable whole serum | | | | | | | | | | | | | | | | |
| Preservative | No preservative added, as it may interfere with the antibody activity. No foreign protein added. | | | | | | | | | | | | | | | | |
| Immunogen | Purified normal IgG isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. | | | | | | | | | | | | | | | | |
| Adsorption | No adsorption required. | | | | | | | | | | | | | | | | |
| Identity & Specificity | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in swine serum, as tested by immunoelectrophoresis and double radial immunodiffusion. | | | | | | | | | | | | | | | | |
| Cross-reactivity | Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has been tested in double radial immunodiffusion with the following results: <table border="0" style="margin-left: 40px;"> <tr> <td>bovine +</td> <td>dog +</td> <td>guinea pig +</td> <td>mouse +</td> <td>rat +</td> </tr> <tr> <td>cat +</td> <td>duck -</td> <td>human +</td> <td>monkey +</td> <td>sheep +</td> </tr> <tr> <td>chicken -</td> <td>goat +</td> <td>horse +</td> <td>pigeon -</td> <td>turkey -</td> </tr> </table> | | bovine + | dog + | guinea pig + | mouse + | rat + | cat + | duck - | human + | monkey + | sheep + | chicken - | goat + | horse + | pigeon - | turkey - |
| bovine + | dog + | guinea pig + | mouse + | rat + | | | | | | | | | | | | | |
| cat + | duck - | human + | monkey + | sheep + | | | | | | | | | | | | | |
| chicken - | goat + | horse + | pigeon - | turkey - | | | | | | | | | | | | | |
| Protein concentration | Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added. | | | | | | | | | | | | | | | | |
| Antibody titre | Precipitin titre not less than 1:64 when tested against normal swine serum in agar block titration. | | | | | | | | | | | | | | | | |
| Intended use | In immunoelectrophoresis and double radial immunodiffusion to identify the presence of IgG; as secondary antibody to precipitate the immunoglobulin in normal swine serum; to prepare an immunoabsorbent for the production of Ig-free swine serum, plasma or non-Ig preparations. | | | | | | | | | | | | | | | | |
| Directions for use | In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells. | | | | | | | | | | | | | | | | |
| Handling | The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. | | | | | | | | | | | | | | | | |
| Packing | Vial with 1 ml lyophilized antiserum. | | | | | | | | | | | | | | | | |
| Storage / shelf life | Lyophilized at +4°C | at least 10 years | | | | | | | | | | | | | | | |
| | reconstituted at or below -20°C | 3-5 years | | | | | | | | | | | | | | | |
| | reconstituted at +4°C | 7 days | | | | | | | | | | | | | | | |
| Caution | This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal and of all associated materials. For <i>in vitro</i> laboratory research purposes only. | | | | | | | | | | | | | | | | |

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl