

<b>Description</b>	<b>Precipitating polyclonal sheep antiserum to bovine serum albumin</b>
<b>Product code</b>	ShAB/Alb
<b>Biological origin</b>	Sheep
<b>Physical form</b>	Delipidated, heat inactivated, lyophilized, stable whole antiserum
<b>Preservative</b>	No preservative added.
<b>Immunogen</b>	Highly purified albumin isolated from bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Adsorption</b>	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.
<b>Identity &amp; Specificity</b>	The defined antibody specificity is directed to albumin as tested against bovine sera. In immunoelectrophoresis and double radial immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the immunogen, a single characteristic precipitin line is obtained which shows a reaction of identity with the precipitin lines obtained against bovine serum and the purified albumin.
<b>Cross-reactivity</b>	Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic determinants. The degree of cross-reactivity is also dependent on the concentrations of the reactants and the sensitivity of the assay arrangement. This antiserum has been tested for cross-reactivity in radial immunodiffusion with the following results: cat - duck - hamster - monkey - rabbit - chicken - goat - horse - mouse - rat + dog - guinea pig - human - pigeon - swine ±
<b>Protein concentration</b>	Total protein and IgG concentrations in the antiserum are comparable to those in pooled sheep serum. No foreign proteins added.
<b>Antibody titre</b>	Precipitin titre not less than 1:64 when tested against normal bovine serum in agar block immunodiffusion titration.
<b>Intended use</b>	As precipitating antiserum to identify or measure bovine albumin by a variety of immunodiffusion techniques, including immunoelectrophoresis, single and double radial immunodiffusion (Mancini, Ouchterlony) and electroimmunodiffusion (Laurell). It has not been tested for use in nephelometry, ELISA or immunochemistry, but this does not exclude such use if proper controls are included.
<b>Directions for use</b>	In immunoelectrophoresis use 2 µl serum, plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single radial immunodiffusion and electroimmunodiffusion use 0.5 to 1.0 percent antiserum in the agar gel.
<b>Handling</b>	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.
<b>Packing</b>	Vial with 1 ml lyophilized antiserum.
<b>Storage / shelf life</b>	Lyophilized at +4°C at least 10 years reconstituted at or below -20°C 3-5 years reconstituted at +4°C 7 days
<b>Caution</b>	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.

**NORDIC IMMUNOLOGICAL LABORATORIES**  
Langendijk 25, 5652 AX Eindhoven, The Netherlands  
Tel. +31 630 070 625, Fax: +31 402 920 069  
E-mail: info@nordiclabs.nl  
www.nordiclabs.nl