

Description	Precipitating polyclonal sheep antiserum to human factor I	
Product code	ShAHu/FI	
Biological origin	Sheep	
Physical form	Delipidated, heat inactivated, lyophilized, stable IgG fraction.	
Preservative	No preservative added	
Immunogen	<p>Factor I is a serine protease composed of two disulphide-linked chains. The protein has a molecular weight of 88.000 and its mean concentration in human plasma is 35 µg/ml. It is involved in the regulation of the C3/C5 convertase of either pathway. Using C4-binding protein or factor H splits the a chains of C4b or C3b respectively, which includes their roles in the C3/C5 convertases. It has been isolated as a homogenous protein for use in the antiserum production.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies reacting with other human serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Identity & Specificity	The reactivity of the antiserum is restricted to factor I. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against human EDTA plasma a single precipitin line is obtained and shows a reaction of full identity with the purified immunogen.	
Cross-reactivity	The antiserum does not cross-react with any other human plasma proteins as tested in gel-diffusion techniques. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins, since homologous proteins of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.	
Protein concentration	Total protein and IgG concentration are comparable to those of pooled normal sheep serum. No foreign proteins added.	
Antibody titre	Precipitin titre not less than 1:128 when tested against normal human plasma in agar-block immunodiffusion titration.	
Intended use	In precipitating techniques as immunoelectrophoresis and single and double radial immunodiffusion (Mancini, Ouchterlony) to identify the presence of F I or to determine its concentration. The presence of non-precipitating antibodies has not been assayed. This does not exclude the use of the antiserum in non-precipitating antibody-binding techniques if proper controls are included.	
Directions for use	In immunoelectrophoresis in agarose-plates use 2 µl human plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl plasma samples (neat and serially diluted) in 2 mm diameter peripheral wells. In electroimmunodiffusion (Laurell) use a concentration of 1% of the antiserum in the gel.	
Packing	Vial with 1 ml lyophilized antiserum.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Handling	<p>The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum.</p> <p>Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.</p>	
Caution	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl