Description	**Precipitating polyclonal sheep antiserum to rabbit IgM, Fc specific**
Product code	ShAR/IgM(Fc)
Biological origin	Sheep
Physical form	Delipidated, heat inactivated, lyophilized, stable whole antiserum
Preservative	No preservative added
Immunogen	Highly purified normal IgM isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Adsorption	Adsorbed using antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins.
Identity & Specificity	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal rabbit serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgM. No precipitation reaction is obtained with purified IgG, IgA, and IgG/Fab fragments.
Cross-reactivity	The antiserum does not cross-react with any other component of the rabbit Ig system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.
Protein concentration	Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.
Antibody titre	Precipitin titre 1:64 when tested against pooled normal rabbit serum in agar-block immunodiffusion titration.
Intended use	In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgM in rabbit serum and other body fluids or to determine its concentration. To prepare an immunoadsorbent for the purification of rabbit IgM from serum or plasma.
Directions for use	In immunoelectrophoresis use 2 μl or equivalent against 120 μl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μl in a 3 mm diameter center well and 2 μl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.
Handling	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.
Packing	Vial with 1 ml lyophilized antiserum.
Storage / shelf life	Lyophilized at +4°C — at least 10 years reconstituted at or below -20°C — 3-5 years reconstituted at +4°C — 7 days
Caution	This antiserum should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For *in vitro* laboratory research purposes only.

NORDIC IMMUNOLOGICAL LABORATORIES
Langendijk 25, 5652 AX Eindhoven, The Netherlands
Tel. +31 630 070 625, Fax: +31 402 920 069
E-mail: info@nordiclabs.nl
www.nordiclabs.nl

© Nordic Immunological Laboratories 2004