

Description	Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal swine antiserum to human IgA, Fc specific	
Product code	SwAHu/IgA(Fc)/FITC	
Biological origin	Swine	
Physical form	FITC-coupled purified hyperimmune swine IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).	
Preservative	No preservative added, as it may interfere with the antibody activity.	
Immunogen	Pools of purified IgA isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.	
Purification	Hyperimmune antisera with strong precipitating activity are selected for fractionation and purification of the IgG (7S) fraction containing the bulk of the defined antibody specificity. It is free of other serum proteins as tested by immunoelectrophoresis.	
Adsorption	Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.	
Identity & Specificity	The reactivity of the antiserum is directed to the Fc subunit of the IgA molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial immunodiffusion using various antiserum concentrations against human serum, a single precipitin line has been obtained which shows a reaction of identity with the precipitin lines obtained with the purified IgA used as immunogens. It does not react with IgG, IgG/Fab fragments and IgM or any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion.	
Cross-reactivity	This immunoconjugate is not species-specific since inter-species cross-reactivity is a normal feature of antisera to immunoglobulins. Cross-reactivity of this antiserum has not been tested in detail.	
Physicochemical characteristics	IgG protein concentration 10 mg/ml. Fluorochrome/IgG protein molar ratio (F/P) is approximately 2.3. No foreign proteins added.	
Fluorescent marker	Fluorescein isothiocyanate isomer 1. Excitation: 492 nm, emission: 515 nm.	
Conjugation procedure	A proprietary technique for the binding of FITC is used, followed by several purification steps to remove free reactants and protein aggregates. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilized to assure stability and long shelf life.	
Intended use	<p>In immunocytochemical and immunohistochemical staining of IgA at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating IgA antibodies in sero-diagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of human origin known to be of the IgA isotype in the middle layer of the indirect test procedure. Antisera to IgA do not discriminate between serum IgA (monomeric and dimeric) and higher molecular forms such as secretory IgA.</p> <p><i>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</i></p> <p>Working dilutions are usually between 1:20 and 1:80, depending on the method used.</p>	
Handling	The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.	
Packing	Vial with 1 ml lyophilized immunoconjugate.	
Storage / shelf life	Lyophilized at +4°C	at least 10 years
	reconstituted at or below -20°C	3-5 years
	reconstituted at +4°C	7 days
Caution	This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For <i>in vitro</i> laboratory research purposes only.	

NORDIC IMMUNOLOGICAL LABORATORIES
 Langendijk 25, 5652 AX Eindhoven, The Netherlands
 Tel. +31 630 070 625, Fax: +31 402 920 069
 E-mail: info@nordiclabs.nl
 www.nordiclabs.nl